

Interacting Phantom Energy and Avoidance of the Big Rip Singularity

Ruben Curbelo, Tame Gonzalez and Israel Quiros*
Universidad Central de Las Villas. Santa Clara. Cuba

(Dated: February 2, 2008)

Models of the universe with arbitrary (non gravitational) interaction between the components of the cosmic fluid: the phantom energy and the background, are investigated. A general form of the interaction that is inspired in scalar-tensor theories of gravity is considered. No specific model for the phantom fluid is assumed. We concentrate our investigation on solutions that are free of the coincidence problem. We found a wide region in the parameter space where the solutions are free of the big rip singularity also. Physical arguments, together with arguments based on the analysis of the observational evidence, suggest that phantom models without big rip singularity might be preferred by Nature.

PACS numbers: 04.20.Jb, 04.20.Dw, 98.80.-k, 98.80.Es, 95.30.Sf, 95.35.+d

I. INTRODUCTION

Recently it has been argued that astrophysical observations might favor a dark energy (DE) component with "supernegative" equation-of-state (EOS) parameter $\omega_i = p_i/\rho_i < -1$ [1, 2], where p_i is the pressure and ρ_i the energy density of the i -th component of the cosmic fluid. Sources sharing this property violate the null dominant energy condition (NDEC). Otherwise, well-behaved energy sources (positive energy density) that violate NDEC (necessarily negative pressure), have EOS parameter less than minus unity [3]. NDEC-violating sources are being investigated as possible dark energy (DE) candidates and have been called as "phantom" components [4, 5]. [27] Since NDEC prevents instability of the vacuum or propagation of energy outside the light cone, then, phantom models are intrinsically unstable. Nevertheless, if thought of as effective field theories (valid up to a given momentum cutoff), these models could be phenomenologically viable [3]. Another very strange property of phantom universes is that their entropy is negative[6].

To the number of unwanted properties of a phantom component with "supernegative" EOS parameter $\omega_{ph} = p_{ph}/\rho_{ph} < -1$, we add the fact that its energy density ρ_{ph} increases in an expanding universe.[28] This property ultimately leads to a catastrophic (future) big rip singularity[8] that is characterized by divergences in the scale factor a , the Hubble parameter H and its time-derivative $\dot{H}$ [10]. Although other kinds of singularity might occur in scenarios with phantom energy, in the present paper we are interested only in the big rip kind of singularity.[29] This singularity is at a finite

amount of proper time into the future but, before it is reached, the phantom energy will rip apart all bound structures, including molecules, atoms and nuclei. To avoid this catastrophic event (also called "cosmic doomsday"), some models and/or mechanisms have been invoked. In Ref.[12], for instance, it has been shown that this singularity in the future of the cosmic evolution might be avoided or, at least, made milder if quantum effects are taken into consideration. Instead, a suitable perturbation of de Sitter equation of state can also lead to classical evolution free of the big rip [13]. Gravitational back reaction effects[14] and scalar fields with negative kinetic energy term with self interaction potentials with maxima[3, 15], have also been considered.

Another way to avoid the unwanted big rip singularity is to allow for a suitable interaction between the phantom energy and the background (DM) fluid[16, 17]. [30] In effect, if there is transfer of energy from the phantom component to the background fluid, it is possible to arrange the free parameters of the model, in such a way that the energy densities of both components decrease with time, thus avoiding the big rip[16]. Models with interaction between the phantom and the DM components are also appealing since the coincidence problem (why the energy densities of dark matter and dark energy are of the same order precisely at present?) can be solved or, at least, smoothed out[16, 17, 20].

Aim of the present paper is, precisely, to study models with interaction between the phantom (DE) and background components of the cosmological fluid but, un-

*Electronic address: rcurbelo@uclv.edu.cu;tame@uclv.edu.cu;israel@uclv.edu.cu

[27] That NDEC-violating sources can occur has been argued decades ago, for instance, in reference [9].

[28] Alternatives to phantom models to account for supernegative EOS parameter have been considered also. See, for instance, references [7]

[29] A detailed study of the kinds of singularity might occur in phantom scenarios (including the big rip) has been the target of ref-

erence [11].

[30] Although experimental tests in the solar system impose severe restrictions on the possibility of non-minimal coupling between the DE and ordinary matter fluids [18], due to the unknown nature of the DM as part of the background, it is possible to have additional (non gravitational) interactions between the dark energy component and dark matter, without conflict with the experimental data. It should be pointed out, however, that when the stability of DE potentials in quintessence models is considered, the coupling dark matter-dark energy is troublesome [19]. Perhaps, the argument might be extended to phantom models.

like the phenomenological approach followed in other cases to specify the interaction term (see references [16, 17]), we start with a general form of the interaction that is inspired in scalar-tensor (ST) theories of gravity. We concentrate our investigation on solutions that are free of the coincidence problem. In this paper no specific model for the phantom energy is assumed. A flat Friedmann-Robertson-Walker (FRW) universe ($ds^2 = -dt^2 + a(t)^2 \delta_{ij} dx^i dx^j$; $i, j = 1, 2, 3$) is considered that is filled with a mixture of two interacting fluids: the background (mainly DM) with a linear equation of state $p_m = \omega_m \rho_m$, $\omega_m = \text{const.}$, [31] and the phantom fluid with EOS parameter $\omega_{ph} = p_{ph}/\rho_{ph} < -1$. Additionally we consider the "ratio function":

$$r = \frac{\rho_m}{\rho_{ph}}, \quad (1)$$

between the energy densities of both fluids. It is, in fact, no more than a useful parametrization. We assume that r can be written as function of the scale factor.

The paper has been organized in the following way: In Section 2 we develop a general formalism that is adequate to study the kind of coupling between phantom energy and the background fluid (inspired in ST theories as said) we want to investigate. Models of references [16, 17] can be treated as particular cases. The way one can escape one of the most undesirable features of any DE model: the coincidence problem, is discussed in Section 3 for the case of interest, where the DE is modelled by a phantom fluid. In this section it is also briefly discussed under which conditions the big rip singularity might be avoided. With the help of the formalism developed before, in Section 4, particular models with interaction between the components of the mixture are investigated where the coincidence problem is solved. The cases with constant and dynamical EOS parameter are studied separately. It is found that, in all cases, there is a wide range in parameter space, where the solutions are free of the big rip singularity also. It is argued that observational data seems to favor phantom models without big rip singularity. This conclusion is reinforced by a physical argument related to the possibility that the interaction term is always bounded. Conclusions are given in Section 5.

II. GENERAL FORMALISM

Since there is exchange of energy between the phantom and the background fluids, the energy is not conserved separately for each component. Instead, the continuity

equation for each fluid shows a source (interaction) term:

$$\dot{\rho}_m + 3H(\rho_m + p_m) = Q, \quad (2)$$

$$\dot{\rho}_{ph} + 3H(\rho_{ph} + p_{ph}) = -Q, \quad (3)$$

where the dot accounts for derivative with respect to the cosmic time and Q is the interaction term. Note that the total energy density $\rho_T = \rho_m + \rho_{ph}$ ($p_T = p_m + p_{ph}$) is indeed conserved: $\dot{\rho}_T + 3H(\rho_T + p_T) = 0$. To specify the general form of the interaction term, we look at a scalar-tensor theory of gravity where the matter degrees of freedom and the scalar field are coupled in the action through the scalar-tensor metric $\chi(\phi)^{-1} g_{ab}$ [22]:

$$S_{ST} = \int d^4x \sqrt{|g|} \left\{ \frac{R}{2} - \frac{1}{2} (\nabla\phi)^2 + \chi(\phi)^{-2} \mathcal{L}_m(\mu, \nabla\mu, \chi^{-1} g_{ab}) \right\}, \quad (4)$$

where $\chi(\phi)^{-2}$ is the coupling function, $\mathcal{L}_m$ is the matter Lagrangian and μ is the collective name for the matter degrees of freedom. It can be shown that, in terms of the coupling function $\chi(\phi)$, the interaction term Q in equations (2) and (3), can be written in the following form:

$$Q = \rho_m H \left[a \frac{d(\ln \bar{\chi})}{da} \right], \quad (5)$$

where we have introduced the following "reduced" notation: $\bar{\chi}(a) \equiv \chi(a)^{(3\omega_m - 1)/2}$ and it has been assumed that the coupling can be written as a function of the scale factor $\bar{\chi} = \bar{\chi}(a)$. [32] This is the general form of the interaction we consider in the present paper. [33] Comparing this with other interaction terms in the bibliography, one can obtain the functional form of the coupling function $\bar{\chi}$ in each case. In Ref. [16], for instance, $Q = 3Hc^2(\rho_{ph} + \rho_m) = 3c^2 H \rho_m (r + 1)/r$, where c^2 denotes the transfer strength. If one compares this expression with (5) one obtains the following coupling function:

$$\bar{\chi}(a) = \bar{\chi}_0 e^{3 \int \frac{da}{a} \left(\frac{r+1}{r} \right) c^2}, \quad (6)$$

where $\bar{\chi}_0$ is an arbitrary integration constant. If $c^2 = c_0^2 = \text{const.}$ and $r = r_0 = \text{const.}$, then $\bar{\chi} =$

[31] Although baryons, which should be left uncoupled or weakly coupled to be consistent with observations, are a non vanishing but small part of the background, these are not being considered here for sake of simplicity. Also there are suggestive arguments showing that observational bounds on the "fifth" force do not necessarily imply decoupling of baryons from DE [21].

[32] If S_{ST} represents Brans-Dicke theory formulated in the Einstein frame, then the coupling function $\chi(\phi) = \chi_0 \exp(-\phi/\sqrt{\omega+3/2})$. The dynamics of such a theory but, for a standard scalar field with exponential self-interaction potential has been studied, for instance, in Ref. [23]. For a phantom scalar field it has been studied in [24].

[33] We recall that we do not consider any specific model for the phantom field so, the scalar-tensor theory given by the action S_{ST} serves just as inspiration for specifying the general form of the interaction term Q .

$\bar{\chi}_0 a^{3e_0^2(r_0+1)/r_0}$. Another example is the interaction term in Ref. [17]: $Q = \delta H \rho_m$, where δ is a dimensionless coupling function. It is related with the coupling function $\bar{\chi}$ (Eqn. (5)) through:

$$\bar{\chi}(a) = \bar{\chi}_0 e^{\int \frac{da}{a} \delta}, \quad (7)$$

and, for $\delta = \delta_0$; $\bar{\chi}(a) = \bar{\chi}_0 a^{\delta_0}$.

If one substitutes (5) in (2), then the last equation can be integrated:

$$\rho_m = \rho_{m,0} a^{-3(\omega_m+1)} \bar{\chi}, \quad (8)$$

where $\rho_{m,0}$ is an arbitrary integration constant.

If one considers Eqn. (1) then, Eqn.(3) can be arranged in the form: $\dot{\rho}_{ph}/\rho_{ph} + 3(\omega_{ph} - 1)H = (3\omega_m - 1)rH[ad(\ln \chi^{-1/2})/da]$ or, after integrating:

$$\rho_{ph} = \rho_{ph,0} \exp\left\{-\int \frac{da}{a} [1 + 3\omega_{ph} + ra \frac{d(\ln \bar{\chi})}{da}]\right\}, \quad (9)$$

where $\rho_{ph,0}$ is another integration constant. Using equations (8), (1) and (9), it can be obtained an equation relating the coupling function $\bar{\chi}$, the phantom EOS parameter ω_{ph} and the "ratio function" r :

$$\bar{\chi}(a) = \bar{\chi}_0 \left(\frac{r}{r+1}\right) e^{-3 \int \frac{da}{a} (\frac{\omega_{ph}-\omega_m}{r+1})}, \quad (10)$$

where, as before, $\bar{\chi}_0$ is an integration constant. Therefore, by the knowledge of $\omega_{ph} = \omega_{ph}(a)$ and $r = r(a)$, one can describe the dynamics of the model under study. Actually, if ω_{ph} and r are given as functions of the scale factor, then one can integrate in Eqn. (10) to obtain $\bar{\chi} = \bar{\chi}(a)$ and, consequently, $\rho_m = \rho_m(a)$ is given through Eqn. (8). The energy density of the phantom field can be computed through either relationship (1) or (9). Besides, the Friedmann equation can be rewritten in terms of r and one of the energy densities:

$$3H^2 = \rho_{ph}(1+r) = \rho_m \left(\frac{1+r}{r}\right), \quad (11)$$

so, the Hubble parameter $H = H(a)$ is also known. Due to Eqn.(11), the (dimensionless) density parameters $\Omega_i = \rho_i/3H^2$ ($\Omega_m + \Omega_{ph} = 1$) are given in terms of only r :

$$\Omega_{ph} = \frac{1}{1+r}, \quad \Omega_m = \frac{r}{1+r}, \quad (12)$$

respectively. It is useful to rewrite Eqn.(10), alternatively, in terms of Ω_{ph} and ω_{ph} :

$$\bar{\chi}(a) = \bar{\chi}_0 (1 - \Omega_{ph}) e^{-3 \int \frac{da}{a} (\omega_{ph} - \omega_m) \Omega_{ph}}. \quad (13)$$

Another useful parameter (to judge whether or not the expansion is accelerated), is the deceleration parameter $q = -(1 + \ddot{H}/H^2)$, that is given by the following equation:

$$q = -1 + \frac{3}{2} \left[\frac{\omega_{ph} + 1 + (\omega_m + 1)r}{1 + r} \right], \quad (14)$$

or, in terms of Ω_{ph} and ω_{ph} :

$$q = \frac{1}{2} [1 + 3\omega_m + 3(\omega_{ph} - \omega_m)\Omega_{ph}]. \quad (15)$$

When the EOS parameter of the phantom field is a constant $\omega_{ph} = -(1 + \xi^2) < -1$, $\xi^2 \in R^+$, we need to specify only the behavior of the ratio function $r = r(a)$ or, equivalently, of the phantom density parameter $\Omega_{ph} = \Omega_{ph}(a)$. In this case the equation for determining of the coupling function $\bar{\chi}$ (10) or, alternatively, (13), can be written in either form:

$$\bar{\chi}(a) = \bar{\chi}_0 \left(\frac{r}{r+1}\right) e^{3(1+\xi^2+\omega_m) \int \frac{da}{a(r+1)}}, \quad (16)$$

or,

$$\bar{\chi}(a) = \bar{\chi}_0 (1 - \Omega_{ph}) e^{3(1+\xi^2+\omega_m) \int \frac{da}{a} \Omega_{ph}}. \quad (17)$$

The deceleration parameter is given, in this case, by $q = -1 + (3/2)[(r - \xi^2)/(r + 1)]$ so, to obtain accelerated expansion $r < 2 + 3\xi^2$.

III. AVOIDING THE COINCIDENCE PROBLEM

In this section we discuss the way in which one of the most undesirable features of any DE model (including the phantom) can be avoided or evaded. For completeness we briefly discuss also the conditions under which the doomsday event can be evaded.

A. How to avoid the coincidence problem?

It is interesting to discuss, in the general case, under which conditions the coincidence problem might be avoided in models with interaction among the components in the mixture. In this sense one expects that a regime with simultaneous non zero values of the density parameters of the interacting components is a critical point of the corresponding dynamical system, so the system lives in this state for a sufficiently long period of time and, hence, the coincidence does not arise.

In the present case where we have a two-component fluid: DM plus DE, one should look for models where the dimensionless energy density of the background (DM) and of the DE are simultaneously non vanishing during a (cosmologically) long period of time. Otherwise, we are interested in ratio functions $r(a)$ that approach a constant value $r_0 \sim 1$ at late times. A precedent of this idea but for a phenomenologically chosen interaction term can be found, for instance, in Ref.[16], where a scalar field model of phantom energy has been explored (a self-interacting scalar field with negative kinetic energy term $\rho_{ph} = -\dot{\phi}^2/2 + V(\phi)$). In this case, as already noted, the interaction term is of the form $Q = 3Hc^2(\rho_{ph} + \rho_m) = 3c^2H\rho_m(r+1)/r$. For this kind of interaction, stability of models with constant ratio $r_0 = \rho_m/\rho_{ph}$ (and constant EOS parameter) yields to

very interesting results. In effect, it has been shown in [16] that, if $r_0 < 1$ (phantom-dominated scaling solution), the model is stable. The solution is unstable whenever $r_0 > 1$ (matter-dominated scaling solution). One can trace the evolution of the model that evolves from the unstable (matter-dominated) scaling solution to the stable (phantom-dominated) one. As a consequence, this kind of interaction indicates a phenomenological solution to the coincidence problem [16]. Actually, once the universe reaches the stable phantom-dominated state with constant ratio $\rho_m/\rho_{ph} = \Omega_m/\Omega_{ph} = r_0$, it will live in this state for a very long time. Then, it is not a coincidence to live in this long-living state where, since $r_0 \sim 1$, $\rho_m \sim \rho_{ph}$.

The above stability study can be extended to situations where no specific model for the phantom energy is assumed (like in the case of interest in the present paper) and where a general interaction term Q is considered (in our case Q is given in Eqn.(5)), if one introduces the following dimensionless quantities[11]:

$$x \equiv \frac{\rho_K}{3H^2}, \quad y \equiv \frac{\rho_V}{3H^2}, \quad (18)$$

where the "kinetic" $\rho_K \equiv (\rho_{ph} + p_{ph})/2$ and "potential" $\rho_V \equiv (\rho_{ph} - p_{ph})/2$ terms have been considered. Besides, if one were to consider FRW universe with spatial curvature k , an additional variable:

$$z \equiv -\frac{k}{a^2 H^2}, \quad (19)$$

is needed. The governing equations can be written as the three-dimensional autonomous system:

$$\begin{aligned} x' &= -(1 + \frac{dp_{ph}}{d\rho_{ph}})(3 + \frac{Q}{2H\rho_K})x \\ &\quad + 3x[2x + \frac{2}{3}z + (\omega_m + 1)(1 - x - y - z)], \\ y' &= -(1 - \frac{dp_{ph}}{d\rho_{ph}})[3x + \frac{Q}{2H\rho_V}y] \\ &\quad + 3y[2x + \frac{2}{3}z + (\omega_m + 1)(1 - x - y - z)], \\ z' &= -2z + 3z[2x + \frac{2}{3}z + (\omega_m + 1) \\ &\quad (1 - x - y - z)] \end{aligned} \quad (20)$$

together with the constrain $\Omega_m = 1 - x - y - z$, where the comma denotes derivative with respect to the variable $N \equiv \ln a$. The parameter range is restricted to be $0 \leq x + y + z \leq 1$. In the remaining part of this section, for simplicity, we assume that $\omega_m = 0$, i. e., the background fluid is dust.[34]




FIG. 1: The phase portraits $(x, y) = (\frac{\rho_K}{3H^2}, \frac{\rho_V}{3H^2})$ for the flat space model with constant ratio r_0 and EOS parameter $\omega_{ph,0} = -(1 + \xi^2)$ are shown for background dust ($\omega_m = 0$). In the upper part of the figure we have considered $r_0 < \xi^2$ ($r_0 = 1, \xi^2 = 3$). The point $(-3/2, 5/2)$ is an attractor. In the lower part of the figure $r_0 > \xi^2$ ($r_0 = 1, \xi^2 = 1/3$). All trajectories in phase space diverge from an unstable node (third critical point $(x, y) = (-1/12, 7/12)$). The first critical point is a saddle. In any case the model has a late time phantom dominated attractor solution (the second critical point $(x, y) = (-1/6, 7/6)$).

1. Flat FRW ($k = 0$)

Let us consider first the spatially flat case $k = 0 \Rightarrow z = 0$. Consider, besides, a constant ratio $r = r_0$ and a constant EOS parameter $\omega_{ph} = \omega_{ph,0} = -(1 + \xi^2) < -1$. According to (16), one obtains a coupling function of the form: $\bar{\chi}(a) = \bar{\chi}_0 (r_0/(r_0 + 1)) a^{3(1+\xi^2)/(r_0+1)}$. If one substitutes this expression for the coupling function into Eqn. (5), then:

$$Q = \frac{3(1 + \xi^2)}{r_0 + 1} H \rho_m. \quad (21)$$

The fixed points of the system (20) can be found once the interaction function Q given in Eqn.(21) and the expression $dp_{ph}/d\rho_{ph} = \omega_{ph,0} = -(1 + \xi^2)$ are substituted into (20). The fixed points of the system $(x', y') = (0, 0)$ are: 1) $(x, y) = (0, 1)$, 2) $(x, y) = (-\xi^2/2, 1 + \xi^2/2)$ and 3) $(x, y) = (-\xi^2/[2(r_0 + 1)], (2 + \xi^2)/[2(r_0 + 1)])$. Of these, the first point is a saddle if $r_0 > \xi^2$ and an unstable node if $r_0 < \xi^2$. The second critical point is always a stable node. In this case $\Omega_{ph} = x + y = 1$ so, there is phantom

[34] The details of the stability study will be given in a separate publication, where other kinds of interaction are considered also.

(dark energy) dominance. The third point is an unstable node whenever $r_0 > \xi^2$ and a saddle otherwise. In this case $\Omega_{ph} = 1/(r_0+1)$ and $\Omega_m = r_0/(r_0+1)$ are simultaneously non vanishing so, it corresponds to matter-scaling solution. In Fig. 1 the convergence of different initial conditions to the attractor (phantom dominated) solution is shown in the phase space (x, y) (flat space case) for given values of the parameters r_0 and ξ^2 . If $r_0 > \xi^2$ all trajectories in phase space diverge from an unstable node (third critical point). The first critical point is a saddle. In any case the model has a late-time phantom dominated attractor solution (the second critical point).

2. FRW with spatial curvature ($k \neq 0$)

The critical points ($x' = 0, y' = 0, z' = 0$) are: 1) $(x, y, z) = (0, 1, 0)$ (de Sitter space), 2) $(x, y, z) = (-\xi^2/2, 1 + \xi^2/2, 0)$, 3) $(x, y, z) = (-\xi^2/[2(r_0+1)], (2 + \xi^2)/[2(r_0+1)], 0)$ and 4) $(x, y, z) = (0, 0, 1)$ (Milne space). The first point is always a saddle, meanwhile, the second critical point is always a stable node (a sink). The third critical point is a saddle whenever $r_0 < \xi^2$ and an unstable node (a source) otherwise. This point corresponds to matter-scaling solution where both Ω_m and Ω_{ph} are simultaneously non negligible. The fourth critical point (Milne space) corresponds to curved ($k \neq 0$) vacuum space ($\Omega_m = \Omega_{ph} = 0$). This point is an unstable node if $r_0 < 2 + 3\xi^2$ and a saddle whenever $r_0 > 2 + 3\xi^2$. Note that, even if the evolution of the universe begins with initial conditions close to the fourth critical point (curved flat space), the model drives the subsequent evolution of the universe to the stable node corresponding to a flat universe dominated by the phantom component.

In summary, the way to evade the coincidence problem when an interaction term of the general form (5) is considered, is then clear: One should look for models where, whenever expansion proceeds, both ω_{ph} and the ratio $r = \rho_m/\rho_{ph} = (1/\Omega_{ph}) - 1$, approach constant values $\omega_{ph,0}$ and $r_0 \sim 1$, respectively. Then, if trajectories in phase space move close to the third critical point (matter-scaling solution), these trajectories will stay close to this point for an extended (arbitrarily long) period of time, before subsequently evolving into the phantom dominated solution (second critical point) that is a late-time attractor of the model.[35]

B. How to avoid the big rip?

If we assume expansion (the scale factor grows up) then, to avoid the future big rip singularity, it is necessary that the background fluid energy density ρ_m and the phantom pressure p_{ph} and energy density ρ_{ph} , were

bounded into the future. Once the "ratio function" r (Eqn.(1)) and the parametric EOS for the phantom fluid ($p_{ph} = \omega_{ph}\rho_{ph}$) are considered, the above requirement is translated into the requirement that ρ_{ph} , $\omega_{ph}\rho_{ph}$ and $r\rho_{ph}$ were bounded into the future. Let us consider, for illustration, the case with constant EOS: $\omega_{ph} = \omega_{ph,0} = -(1+\xi^2) < -1$, and the following ratio function: $r = r_0 a^{-k}$; $\xi^2, k \in R^+$ [17]. Integrating in the exponent in Eqn.(16) one obtains the following expression for the coupling function:

$$\bar{\chi}(a) = \bar{\chi}_0 r_0 (a^k + r_0)^{-1+3(1+\xi^2)/k}. \quad (22)$$

The expression for the DM and phantom energy densities are then given through equations (see equations (8) and (1)):

$$\rho_m = \bar{\rho}_{m,0} a^{-3} (a^k + r_0)^{-1+3(1+\xi^2)/k}, \quad (23)$$

and

$$\rho_{ph} = \bar{\rho}_{ph,0} a^{-3+k} (a^k + r_0)^{-1+3(1+\xi^2)/k}, \quad (24)$$

where $\bar{\rho}_{m,0} = \rho_{m,0}\bar{\chi}_0 r_0$ and $\bar{\rho}_{ph,0} = \rho_{m,0}\bar{\chi}_0$. We see that, for $a \gg r_0^{1/k}$, $\rho_m \propto a^{-k+3\xi^2}$, while $\rho_{ph} \propto a^{3\xi^2}$. For $\xi^2 < k/3$ the DM energy density ρ_m decreases with the expansion of the universe. However, the phantom energy density always increases with the expansion and, consequently, the big rip singularity is unavoidable in this example. The conclusion is that, for a phantom fluid with a constant EOS $\omega_{ph} = \omega_{ph,0}$ and a power-law ratio of phantom to background (dust) fluid energy densities $\rho_m/\rho_{ph} \propto a^{-k}$, the big rip singularity is unavoidable.

For a constant ratio $r = r_0$ and a constant EOS ($\omega_{ph} = \omega_{ph,0} = -(1+\xi^2) < -1$) [16], direct integration in Eqn. (16) yields to the following coupling function:

$$\bar{\chi}(a) = \bar{\chi}_0 \left(\frac{r_0}{r_0+1} \right) a^{\frac{3(1+\xi^2)}{r_0+1}}, \quad (25)$$

so, the background and phantom energy densities in Eqn. (8) are given by:

$$\rho_{ph} = r_0^{-1} \rho_m = \frac{\bar{\rho}_{m,0}}{r_0} a^{-3(\frac{r_0-\xi^2}{r_0+1})}, \quad (26)$$

where $\bar{\rho}_{m,0} = \rho_{m,0}\bar{\chi}_0 [r_0/(r_0+1)]$. We see that, whenever there is expansion of the universe, for $r_0 \geq \xi^2$, there is no big rip in the future of the cosmic evolution, since $\rho_{ph} \propto r_0 \rho_{ph} \propto \omega_{ph,0} \rho_{ph} \propto a^{-3(r_0-\xi^2)/(r_0+1)}$ are bounded into the future. In order to get accelerated expansion, one should have $r_0 < 2 + 3\xi^2$ so, the ratio of background (dust) energy density to phantom energy density should be in the range $\xi^2 \leq r_0 < 2 + 3\xi^2$.

IV. MODELS THAT ARE FREE OF THE COINCIDENCE PROBLEM

In the present section we look for a model with an appropriated coupling function $\bar{\chi}$ that makes possible to

[35] This argument is due to Alan Coley (private communication).




FIG. 2: The plot of energy density parameters Ω_m (dashed curve) and Ω_{ph} (thin solid curve) as well as of the ratio function r (thick solid curve) vs redshift z , is shown for the model with constant EOS parameter. For simplicity background dust ($\omega_m = 0$) was considered. The values of the free parameters are: $m = 12$, $C = 0.02$ and $\omega_{ph,0} = -1.1$. The ratio function r approaches a constant value at negative redshift $r_0 = 0.23$. At present ($z = 0$) $r = 0.25$, which means that we are already in the long living matter-scaling state. The background energy density dominates the early stages of the cosmic evolution. At $z \sim 0.33$ both density parameters equate and, since then, the phantom component dominates the dynamics of the expansion. A very similar behavior is obtained for Ω_m , Ω_{ph} and r in the case with dynamical EOS parameter (Case B).

avoid the coincidence problem. We study separately the cases with a constant EOS parameter $\omega_{ph} = \omega_{ph,0} = -(1 + \xi^2) < -1$; $\xi^2 \in R^+$ and with variable EOS parameter.

A. Constant EOS parameter $\omega_{ph} = \omega_{ph,0}$

In order to have models where both Ω_m and Ω_{ph} are simultaneously non-vanishing during a long interval of cosmological time (no coincidence issue) and that, at the same time, are consistent with observational evidence on a matter dominated period with decelerated expansion, lasting until recently (redshift $z \sim 0.39$ [2]), we choose the following dimensionless density parameter for the phantom component:

$$\Omega_{ph} = \frac{m}{B} \left(\frac{a^m}{a^m + C} \right), \quad (27)$$

where m , B , and C are arbitrary constant parameters. This kind of density parameter function can be arranged to fit the relevant observational data, by properly choosing the free parameters (m , B , C) (m controls the curvature of the curve $\Omega_{ph}(z)$, meanwhile C controls the point at which $\Omega_{ph}(z_{eq}) = \Omega_m(z_{eq})$). In this sense, the model under study could serve as an adequate, observationally testable, model of the universe. The choice of Ω_{ph} in

equation (27), when substituted in Eqn. (17) yields to:

$$\bar{\chi}(a) = \bar{\chi}_0 \frac{(1 - \frac{m}{B})a^m + C}{(a^m + C)^{1-3(\omega_m - \omega_{ph,0})/B}}. \quad (28)$$

In consequence, equations (8) and (1) lead to the following expressions determining the background energy density and the energy density of the phantom component respectively;

$$\rho_m = \bar{\rho}_{m,0} a^{-3} \frac{(1 - \frac{m}{B})a^m + C}{(a^m + C)^{1-3(\omega_m - \omega_{ph,0})/B}}, \quad (29)$$

where $\bar{\rho}_{m,0} \equiv \rho_{m,0} \bar{\chi}_0/B$, $\bar{\rho}_0 = \bar{\rho}_{m,0} \bar{\xi}_0$ and

$$\rho_{ph} = \bar{\rho}_{ph,0} a^{m-3} (a^m + C)^{-1+3(\omega_m - \omega_{ph,0})/B}, \quad (30)$$

where $\bar{\rho}_{ph,0} \equiv m \rho_{m,0} \bar{\chi}_0/B$.

It is worth noting that, if $\omega_m - B/m \leq \omega_{ph,0} (< -1)$, there is no big rip singularity in the future of the cosmic evolution as described by the present model. In effect, for large $a \gg 1$, the following magnitudes

$$\begin{aligned} \rho_{ph} &\propto r \rho_{ph} \propto \omega_{ph,0} \rho_{ph} \\ &\propto a^{-3(1-m(\omega_m - \omega_{ph,0})/B)}, \end{aligned} \quad (31)$$

are bounded into the future. In particular, the phantom energy density is a decreasing function of the scale factor. Note that the ratio function $r = \Omega_m/\Omega_{ph}$ approaches the constant value $(B/m) - 1$ so both Ω_m and Ω_{ph} are simultaneously non vanishing. As explained in the former section, this means that the universe is in the third critical point (matter-scaling solution) which is a saddle for $r_0 < \xi^2$. In consequence, the universe will evolve close to this critical point for a sufficiently long period of time and then, the model approaches a stable dark energy dominated regime (second critical point).

In Fig. 2 we show the behavior of the density parameters Ω_m and Ω_{ph} and of the ratio function r as functions of the redshift (for simplicity we have considered background dust: $\omega_m = 0$). The following values of the free parameters: $m = 12$, $\omega_{ph,0} = -1.1$, $C = 0.02$ have been chosen, so that it is a big rip free solution. Besides we set the DE density parameter at present epoch $\Omega_{ph}(z = 0) = 0.8$ so, the following relationship between the free constant parameters B , m and C should be valid: $B = 1.25[m/(1 + C)]$. In consequence, for the values of m and C given above: $B/m = 1.22$ so, $\omega_{ph,0} = -1.1 > -1.22$, meaning that there is no big rip singularity in the future of the cosmic evolution in this model (see the condition $-B/m \leq \omega_{ph,0}$ under Eqn. (30)). In Fig. 3, the plot of the deceleration parameter q vs redshift is shown for three different values of the EOS parameter $\omega_{ph,0} = -1.1$, -1.5 and -3 respectively. It is a nice result that the solution without big rip (thick solid curve) is preferred by the observational evidence, since, according to a model-independent analysis of SNIa data[25], the mean value of the present value of the deceleration parameter is $\langle q_0 \rangle \approx -0.76$.




FIG. 3: The deceleration parameter q is plotted as function of the redshift z for the Model A (constant EOS parameter), for three different values of $\omega_{ph,0} = -1.1, -1.5$ and -3 (thick solid, thin solid and dashed curves respectively). Note that the transition from decelerated (positive q) into accelerated (negative q) expansion occurs at $z \simeq 0.4$ for the value $\omega_{ph,0} = -1.1$. It is apparent that this is, precisely, the observationally favored solution, since, according to a model-independent analysis of SNIa data, $q(z=0) \approx -0.76$ [25]. In consequence, solutions without big rip are preferred by observations.

The way the coincidence problem is avoided in this model is clear from Fig.2 also. The ratio between the energy density parameters of the background and of the DE approaches a constant value $r_0 = 0.23$ for negative redshift so, both Ω_m and Ω_{ph} are non negligible. As already said, this is a critical point (third critical point in the stability study of section 3) and the universe can live for a long time in this state. Moreover, since at present ($z=0$); $r(z=0) = 0.25$, we can conclude that the universe is already in this long living state.

B. Dynamical EOS parameter

We now consider a dynamical EOS parameter $\omega_{ph} = \omega_{ph}(a)$. In this case we should give as input the functions $\Omega_{ph}(a)$ and $\omega_{ph}(a)$. In consequence, one could integrate in the exponent in Eqn.(13) so that the dynamics of the model is completely specified. In order to assure avoiding of the coincidence problem, let us consider the same phantom energy density parameter as in the former subsection (Eqn. (27)), but rewritten in a simpler form:

$$\Omega_{ph}(a) = \frac{\alpha a^m}{a^m + \beta}, \quad (32)$$

where m, α and β are non negative (constant) free parameters. To choose the function $\omega_{ph}(a)$ one should take into account the following facts: i) at high redshift and until recently ($z \simeq 0.39 \pm 0.03$ [2]) the expansion was decelerated (positive deceleration parameter q) and, since then, the expansion accelerates (negative q), ii) for negative redshifts the EOS parameter approaches a constant value




FIG. 4: The plot of the energy densities of DM ρ_m (upper part of the figure), phantom component ρ_{ph} (middle part) and of the Hubble parameter H vs redshift, is shown for three values of the constant parameter $\omega_{ph,0}$: -1.1 (thick solid curve), -1.5 (thin solid curve) and -3 (dashed curve) for the model with dynamical EOS parameter. It is apparent that, only in the first case the model is big rip free. Note that, due to the values chosen for the free parameters, in particular $m = 12$, the phantom energy density vanishes as one approaches the initial big bang singularity. This is true whenever $m > 3$ (see Eqn. (36)).

$\omega_{ph,0}$ more negative than -1 ; $\omega_{ph,0} = -(1+\xi^2)$, $\xi^2 \in R^+$. We consider, additionally, the product $(\omega_{ph} - \omega_m) \Omega_{ph}$ to be a not much complex function, so that the integral in the exponent in Eqn.(13) could be taken analytically. A function that fulfills all of the above mentioned requirements is the following:

$$\omega_{ph}(a) = \omega_m + \omega_{ph,0} \frac{(a^m + \beta)(a^m - \delta)}{a^{2m} + \delta}, \quad (33)$$

where δ is another free parameter. The parameter m controls the curvature of the density parameter function




FIG. 5: The interaction term Q is plotted vs redshift z . Note that for $0.2 \lesssim z \lesssim 1.7$, Q is negative, meaning that there is transfer of energy from the DM into DE component. For higher z the model proceeds without interaction. Q is bounded into the future only for the big rip free case (thick solid curve).

$\Omega_{ph}(z)$, while δ controls the point of equality $\Omega_{ph}(z_{eq}) = \Omega_m(z_{eq})$. The resulting coupling function is

$$\bar{\chi}(a) = \bar{\chi}_0 (1 - \Omega_{ph}) (a^{2m} + \delta)^{-\frac{3\alpha}{2m}\omega_{ph,0}} \times e^{\frac{3\alpha\sqrt{\delta}}{m} \arctan(\frac{a^m}{\sqrt{\delta}})}. \quad (34)$$

Therefore, the background and phantom energy densities are given by the following expressions:

$$\rho_m(a) = \rho_{m,0} \left[\frac{(1-\alpha)a^m + \beta}{a^m + \beta} \right] a^{-3(\omega_m+1)} \times (a^{2m} + \delta)^{-\frac{3\alpha}{2m}\omega_{ph,0}} e^{\frac{3\alpha\sqrt{\delta}}{m} \arctan(\frac{a^m}{\sqrt{\delta}})}, \quad (35)$$

$$\rho_{ph}(a) = \rho_{ph,0} \left[\frac{a^{m-3(\omega_m+1)}}{a^m + \beta} \right] (a^{2m} + \delta)^{-\frac{3\alpha}{2m}\omega_{ph,0}} \times e^{\frac{3\alpha\sqrt{\delta}}{m} \arctan(\frac{a^m}{\sqrt{\delta}})}, \quad (36)$$

where the constant $\rho_{ph,0} = \alpha \rho_{m,0}$. At late times (large a), the functions r , $\rho_{ph} \propto \omega_{ph}$, $\rho_{ph} \propto \rho_{ph} \propto a^{-3(\omega_m+1+\alpha\omega_{ph,0})}$. This means that, whenever $-(1 + \omega_m)/\alpha \leq \omega_{ph,0} (< -1)$, these functions are bounded into the future and, consequently, there is no big rip in the future of the cosmic evolution in the model under study.

In Fig.4 the plot of the energy densities of DM ρ_m (upper part of the figure), phantom component ρ_{ph} (middle part) and of the Hubble parameter H vs redshift, is shown for three values of the constant parameter $\omega_{ph,0}$: -1.1 (thick solid curve), -1.5 (thin solid curve) and -3 (dashed curve). The following values of the free parameters: $m = 12$, $\beta = 0.03$, and $\delta = 3 \cdot 10^{-4}$ have been chosen. We assume $\Omega_m(z=0) = 0.3$ [26] so, the following relationship between α and β should take place: $\alpha = 0.7(\beta + 1)$. It is apparent that, only in the first case ($\omega_{ph,0} = -1.2$) the model is big rip free. Actually, as already noted, the necessary requirement for absence of big rip is, for the chosen set of parameters,




FIG. 6: In the upper and middle part of the figure the EOS parameter ω_{ph} and the deceleration parameter q are plotted as functions of the redshift z (Model B), for three values of the constant parameter $\omega_{ph,0} = -1.1, -1.5$ and -3 (thick solid, thin solid and dashed curves respectively). It is worth noting that at high redshift ω_{ph} approaches a constant value $\omega_{ph} = 0$. It is a minimum at $z = 0.35$. The dark energy became a phantom fluid only recently, at $z \sim 0.45$. Note that the transition from decelerated (positive q) into accelerated (negative q) expansion occurs at $z \simeq 0.4$ for the value $\omega_{ph,0} = -1.1$. This last (big rip free) solution, seems to be preferred by observations since, according to Ref. [25], $q(z=0) \approx -0.76$. In the lower part of the figure the ratio function is plotted for three values of the parameter m : 12 (thick solid curve), 8 (thin solid curve) and 4 (dashed curve) respectively. In any case, at present, the universe is already in the state with constant ratio r .

$\omega_{ph,0} \geq -1/\alpha = -1.387$. It is noticeable that, since we chose $m = 12 > 3$, the DE energy density diminishes as one goes back into the past (it is a maximum at present). In all cases the interaction term is negative during a given period of time in the past (see Fig.5), meaning that the DM transferred energy to the DE component.

At higher redshift the evolution proceeds without interaction. Worth noting that only in the singularity free case ($\omega_{ph,0} = -1.2$) the interaction term is bounded into the future.

A physical argument against doomsday event can be based on the following analysis. At late time, the interaction term $Q \simeq -3\alpha \omega_{ph,0} \rho_m H$. This means that, only in the case when the evolution proceeds without big rip (ρ_m and H are bounded), the interaction between the components of the cosmic fluid does not become unphysically large at finite time into the future. Besides, for the big rip free case, as the expansion proceeds into the future, the Hubble parameter approaches a negligible value, meaning that the ratio Q/ρ_m is small.

In the model the transition from decelerated into accelerated expansion takes place at $z \simeq 0.4$ (see Fig. 6). At late times the deceleration parameter approaches the constant value $q_0 \approx -0.8$. Therefore, as in the case with constant EOS parameter (Model A), the observational evidence (as argued in Ref. [25], $q(z=0) \approx -0.76$) seems to favor solutions without big rip singularity. It is interesting to note that, in this model, the dark energy component behaved like dust ($\omega_{ph} = 0$) at early times (high z), i. e., it behaved like an "ordinary" fluid with attractive gravity and, just until very recently ($z \simeq 0.45$), it has not become a phantom fluid. At present, the universe is already in a state characterized by constant $\omega_{ph,0} \sim -1.2$ and $r_0 = 0.48$, meaning that $\Omega_{m,0} = r_0/(r_0 + 1)$ and $\Omega_{ph,0} = 1/(r_0 + 1)$ are simultaneously non vanishing. As shown before, this is a critical point of the corresponding dynamical system so, probably, the universe will stay in this state for a long time and, consequently, the coincidence problem does not arise.

V. CONCLUSIONS

In this paper we have investigated models with additional (non gravitational) interaction between the phantom component and the background. This kind of interaction is justified if the interacting components are of unknown nature, as it is the case for the DM and the DE, the dominant components in the cosmic fluid. For the sake of simplicity, baryons, a non vanishing, but small component of the background, have not been considered in the present study, however, in a more complete study these should be included also. A comment should be made in this regard: There are suggestive arguments showing that observational bounds on the "fifth" force do not necessarily imply decoupling of baryons from DE[21].

Unlike the phenomenological approach followed in other cases to specify the interaction term (see references [16, 17]), we started with a general form of the interaction that is inspired in ST theories of gravity (see Eqn.(5)). We have considered no specific model for the phantom fluid. Within this context, cosmological evolutions without coincidence problem have been the target of this investigation. We have studied two different mod-

els: Model A, where the DE EOS parameter is always a constant (phantom DE) and Model B, where we considered a dynamical DE EOS parameter (the DE became a phantom just recently). In these models there is a wide range in the parameter space where the cosmic evolution is free of the unwanted big rip singularity also.

The interaction between the different components of the cosmic mixture, enables the cosmic evolution to proceed without big rip singularity. In effect, if the interaction term is chosen so that there is transfer of energy from the phantom component to the background, then it is possible to arrange the free parameters of the model to account for decreasing energy density of the phantom fluid (as well as of the background fluid). A physical argument against big rip can be based on the analysis of the interaction term Q . In model B, for instance, at late time the interaction term $Q \simeq -3\alpha \omega_{ph,0} \rho_m H$. This means that, only whenever ρ_m and H are bounded (no big rip), the interaction between the components of the cosmic fluid does not become unphysically large at finite time into the future. Besides, for the big rip free case, as the expansion proceeds into the future, the Hubble parameter approaches a negligible value, meaning that the ratio Q/ρ_m is small. From the observational perspective it seems that the evolution without doomsday event is preferred also. For this purpose we have considered model-independent analysis of SNIa data yielding to a mean value of the present value of the deceleration parameter $q_0 > \approx -0.76$ [25].

Interacting models of dark energy are also useful to account for the coincidence problem. In the present study, it has been shown that, a solution with simultaneously non vanishing values $\Omega_{m,0} = r_0/(r_0 + 1)$ and $\Omega_{ph,0} = 1/(r_0 + 1)$ (r_0 is the constant ratio of the DM to DE densities), is a critical point of the corresponding dynamical system (third critical point in section 3). Hence, if trajectories in phase space get close to this point, these trajectories will stay there during a sufficiently long period of time. This means that, once the universe is driven into the state with simultaneously non vanishing Ω_m and Ω_{ph} , it will live in this state for a (cosmologically) long period of time until it is finally attracted into the phantom dominated phase being a stable node of the models (second critical point in section 3). This is the way the coincidence problem is solved in the present investigation. It seems that we live already in the matter-scaling regime (third critical point) and will live in it for long time until, finally, the phantom dominates the destiny of the cosmic evolution. In any case the evolution of the universe will proceed without the risk of any catastrophic (doomsday) destiny.

An obvious limitation of the present study is that only two specific models (model A and model B) have been considered. However, we think this suffices to illustrate how interacting models can be constructed that avoid the coincidence problem and, at the same time, fit some of the observational evidence. The outcome that big rip free models are preferred by some of these observations

is just a nice result.

We are very grateful to Alan Coley and Sigbjorn Hervik for valuable conversations that helped us to improve the original version of the manuscript. We thank Marc Kamionkowski and Bret McInnes for calling our attention upon references [8] and [13] respectively. We

thank also Joerg Jaeckel, Sergei Odintsov, Vakif Onemli, Diego Pavon and Thanu Padmanabhan for useful comments and for pointing to us important references. The authors thank the MES of Cuba by partial financial support of this research.

-
- [1] R. A. Knop et al., *Astrophys. J.* **598** (2003) 102 (astro-ph/0309368); A. G. Riess et al. [Supernova Search Team Collaboration], *Astrophys. J.* **607** (2004) 665 (astro-ph/0402512).
 - [2] U. Alam, V. Sahni, T. D. Saini and A. A. Starobinsky, *Mon. Not. Roy. Astron. Soc.* **354** (2004) 275 (astro-ph/0311364); U. Alam, V. Sahni and A. A. Starobinski, *JCAP* **0406** (2004) 008 (astro-ph/0403687); B. Feng, X. L. Wang and X. Zhang, *Phys. Lett. B* **607** (2005) 35-41 (astro-ph/0404224); T. R. Choudhury and T. Padmanabhan, *Astron. Astrophys.* **429** (2005) 807 (astro-ph/0311622); H. K. Jassal, J. S. Bagla and T. Padmanabhan, *Mon. Not. Roy. Astron. Soc. Letters* **356** (2005) L11-L16 (astro-ph/0404378).
 - [3] S. M. Carroll, M. Hoffman and M. Trodden, *Phys. Rev. D* **68** (2003) **023509** (astro-ph/0301273).
 - [4] R. R. Caldwell, *Phys. Lett. B* **545** (2002) 23-29 (astro-ph/9908168).
 - [5] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **562** (2003) 147-152 (hep-th/0303117).
 - [6] I. Brevik, S. Nojiri, S. D. Odintsov and L. Vanzo, *Phys. Rev. D* **70** (2004) 043520 (hep-th/0401073).
 - [7] V. K. Onemli and R. P. Woodard, *Class. Quant. Grav.* **19** (2002) 4607 (gr-qc/0204065); *Phys. Rev. D* **70** (2004) 107301 (gr-qc/0406098).
 - [8] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, *Phys. Rev. Lett.* **91** (2003) 071301 (astro-ph/0302506).
 - [9] H. P. Nilles, *Phys. Rept.* **110** (1984) 1; J. D. Barrow, *Nucl. Phys. B* **310** (1988) 743; M. D. Pollock, *Phys. Lett. B* **215** (1988) 635.
 - [10] L. P. Chimento and R. Lazkoz, *Mod. Phys. Lett. A* **19** (2004) 2479-2484 (gr-qc/0405020).
 - [11] S. Nojiri, S. D. Odintsov and S. Tsujikawa, *Phys. Rev. D* **71** (2005) 063004 (hep-th/0501025).
 - [12] S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **595** (2004) 1-8 (hep-th/0405078); *Phys. Rev. D* **70** (2004) 103522 (hep-th/0408170); E. Elizalde, S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **70** (2004) 043539 (hep-th/0405034).
 - [13] B. McInnes, *JHEP* **0208** (2002) 029 (hep-th/0112066).
 - [14] P. Wu and H. Yu, *Nucl. Phys. B* **727** (2005) 355-367 (astro-ph/0407424).
 - [15] P. Singh, M. Sami and N. Dadhich, *Phys. Rev. D* **68** (2003) 023522 (hep-th/0305110).
 - [16] Z. K. Guo and Y. Z. Zhang, *Phys. Rev. D* **71** (2005) 023501 (astro-ph/0411524).
 - [17] R. G. Cai and A. Wang, *JCAP* **0503** (2005) 002 (hep-th/0411025).
 - [18] C. M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, 1993); gr-qc/0103036.
 - [19] M. Doran and J. Jaeckel, *Phys. Rev. D* **66** (2002) 043519 (astro-ph/0203018).
 - [20] W. Zimdahl, D. Pavon and L. P. Chimento, *Phys. Lett. B* **521** (2001) 133-138 (astro-ph/0105479); L. P. Chimento, A. S. Jakubi, D. Pavon and Winfried Zimdahl, *Phys. Rev. D* **67** (2003) 083513 (astro-ph/0303145); L. P. Chimento, A. S. Jakubi and D. Pavon, *Phys. Rev. D* **62** (2000) 063508 (astro-ph/0005070); L. P. Chimento, A. S. Jakubi and D. Pavon, *Phys. Rev. D* **67** (2003) 087302 (astro-ph/0303160).
 - [21] L. P. Chimento, A. S. Jakubi, D. Pavon and W. Zimdahl, *Phys. Rev. D* **67** (2003) 083513 (astro-ph/0303145).
 - [22] N. Kaloper and K. A. Olive, *Phys. Rev. D* **57** (1998) 811-822 (hep-th/9708008).
 - [23] A. P. Billyard and A. A. Coley, *Phys. Rev. D* **61** (2000) 083503 (astro-ph/9908224).
 - [24] Z. K. Guo, R. G. Cai and Y. Z. Zhang, *JCAP* **0505** (2005) 002 (astro-ph/0412624).
 - [25] M. V. John, *Astrophys. J.* **614** (2004) 1 (astro-ph/0406444).
 - [26] P. J. E. Peebles, astro-ph/0410284.